

Connecting the geometric measure of entanglement and entanglement witnesses

Tzu-Chieh Wei and Paul M. Goldbart

Department of Physics, University of Illinois at Urbana-Champaign,
1110 West Green Street, Urbana, Illinois 61801-3080, U.S.A.

(Dated: February 1, 2008)

The geometric measure of entanglement is an approach to quantifying entanglement that is based on the Hilbert-space distance (or, equivalently, angle) between pure states and their best unentangled approximants. An entanglement witness is an operator that reveals entanglement for a given entangled state. A connection is identified between entanglement witnesses and the geometric measure of entanglement. This offers a new interpretation of the geometric measure of entanglement of a state, and renders it experimentally verifiable, doing so most readily for states that are pure.

PACS numbers: 03.67.Mn, 03.65.Ud

Introduction: Entanglement is now recognized as a resource central to much of quantum information processing [1]. Thus, characterizing and quantifying entanglement has emerged as a prominent theme in quantum information theory. Achievements in quantifying mixed-state entanglement lie primarily in bipartite settings [2]. For *multipartite* mixed states the issue of entanglement evidently presents even greater challenges.

Recent research on quantifying multipartite entanglement has explored a geometric approach. First introduced by Shimony [3] in the setting of bipartite pure states, this geometric approach has been generalized to multipartite settings by Barnum and Linden [4], and further developed in Ref. [5]. In the present Paper, our aim is to identify a connection between two apparently distinct aspects of entanglement: entanglement witnesses [6] and the geometric approach to entanglement. As entanglement witnesses are observables, and can, in principle, be measured, the geometric measure of entanglement thus becomes verifiable experimentally.

Geometric measure of entanglement (GME): We begin by briefly reviewing the formulation of the GME in the pure-state setting. Let us start by analyzing a multipartite system comprising n parts, each of which may have a distinct Hilbert space. Consider the general n -partite pure state, expanded in the local bases $\{|e_{p_i}^{(i)}\rangle\}$: $|\psi\rangle = \sum_{p_1 \dots p_n} \chi_{p_1 p_2 \dots p_n} |e_{p_1}^{(1)} e_{p_2}^{(2)} \dots e_{p_n}^{(n)}\rangle$. As shown in Ref. [5], the closest separable pure state

$$|\phi\rangle \equiv \otimes_{i=1}^n |\phi^{(i)}\rangle = \otimes_{i=1}^n \sum_{p_i} c_{p_i}^{(i)} |e_{p_i}^{(i)}\rangle \quad (1)$$

obeys the equations

$$\sum_{p_1 \dots \widehat{p_i} \dots p_n} \chi_{p_1 p_2 \dots p_n}^* c_{p_1}^{(1)} \dots \widehat{c_{p_i}^{(i)}} \dots c_{p_n}^{(n)} = \Lambda c_{p_i}^{(i)*}, \quad (2a)$$

$$\sum_{p_1 \dots \widehat{p_i} \dots p_n} \chi_{p_1 p_2 \dots p_n} c_{p_1}^{(1)*} \dots \widehat{c_{p_i}^{(i)*}} \dots c_{p_n}^{(n)*} = \Lambda c_{p_i}^{(i)}, \quad (2b)$$

where the eigenvalue $\Lambda \in [-1, 1]$ is associated with the Lagrange multiplier enforcing the constraint $\langle \phi | \phi \rangle = 1$, and $\widehat{}$ denotes exclusion. As discussed in Ref. [5], the

spectrum of eigenvalues $\{\Lambda\}$ can be interpreted as the cosine of the angle between $|\psi\rangle$ and stationary states $\{|\phi\rangle\}$. Furthermore, the largest eigenvalue $\Lambda_{\max}$, which we call the *entanglement eigenvalue*, corresponds to the separable state closest to $|\psi\rangle$. An equivalent way to view $\Lambda_{\max}$ is via

$$\begin{aligned} \Lambda_{\max}^2(|\psi\rangle) &= \max_{\text{separable } \phi} ||\langle \phi | \psi \rangle||^2 \\ &= \max_{\text{separable } \phi} \text{Tr}(|\phi\rangle\langle\phi| |\psi\rangle\langle\psi|). \end{aligned} \quad (3)$$

The precise measure of the entanglement of $|\psi\rangle$ adopted in Ref. [5] is $E_{\sin^2} \equiv 1 - \Lambda_{\max}^2$.

Entanglement witness (EW): The entanglement witness $\mathcal{W}$ for an entangled state ρ is defined to be an operator that is Hermitian and obeys the following conditions:

- (i) $\text{Tr}(\mathcal{W}\sigma) \geq 0$ for all separable states σ , and
- (ii) $\text{Tr}(\mathcal{W}\rho) < 0$.

Here, we wish to establish a relationship between $\Lambda_{\max}$ for the entangled pure state $|\psi\rangle$ and the optimal element of the set of entanglement witnesses $\mathcal{W}$ for $|\psi\rangle$ that have the specific form

$$\mathcal{W} = \lambda^2 \mathbb{1} - |\psi\rangle\langle\psi|, \quad (4)$$

this set being parametrized by the real, non-negative number λ^2 . By *optimal* we mean that, for this specific form of witnesses, the value of the “detector” $\text{Tr}(\mathcal{W}|\psi\rangle\langle\psi|)$ is as negative as can be.

In order to satisfy condition (i) we must ensure that, for any *separable* state σ , we have $\text{Tr}(\mathcal{W}\sigma) \geq 0$. As the density matrix for any separable state can be decomposed into a mixture of *separable pure* states [i.e., $\sigma = \sum_i |\phi_i\rangle\langle\phi_i|$ where $\{|\phi_i\rangle\}$ are separable pure states], condition (i) will be satisfied as long as $\text{Tr}(\mathcal{W}|\phi\rangle\langle\phi|) \geq 0$ for all separable *pure* states $|\phi\rangle$. This condition is equivalent to

$$\lambda^2 - ||\langle\psi|\phi\rangle||^2 \geq 0 \quad (\text{for all separable } |\phi\rangle), \quad (5)$$

which leads to

$$\lambda^2 \geq \max_{|\phi\rangle} ||\langle\psi|\phi\rangle||^2 = \Lambda_{\max}^2(|\psi\rangle). \quad (6)$$

Condition (ii) requires that $\text{Tr}(\mathcal{W}|\psi\rangle\langle\psi|) < 0$, in order for $\mathcal{W}$ to be a valid EW for $|\psi\rangle$; this gives $\lambda^2 - 1 < 0$. Thus, we have established the range of λ for which $\lambda^2 \mathbb{1} - |\psi\rangle\langle\psi|$ is a valid EW for $|\psi\rangle$:

$$\Lambda_{\max}^2(|\psi\rangle) \leq \lambda^2 < 1. \quad (7)$$

With these preliminaries in place, we can now establish the connection we have been seeking. Of the specific family (4) of entanglement witnesses for $|\psi\rangle$ that we have been considering, the one of the form $\mathcal{W}_{\text{opt}} = \Lambda_{\max}^2(|\psi\rangle)\mathbb{1} - |\psi\rangle\langle\psi|$ is optimal, in the sense that it achieves the most negative value for the detector $\text{Tr}(\mathcal{W}_{\text{opt}}|\psi\rangle\langle\psi|)$:

$$\min_{\mathcal{W}} \text{Tr}(\mathcal{W}|\psi\rangle\langle\psi|) = \text{Tr}(\mathcal{W}_{\text{opt}}|\psi\rangle\langle\psi|) = -E_{\sin^2}(|\psi\rangle), \quad (8)$$

where $\mathcal{W}$ runs over the class (4) of witnesses.

Some illustrative examples: For the state $|\text{GHZ}\rangle \equiv (|000\rangle + |111\rangle)/\sqrt{2}$ the optimal witness is

$$\mathcal{W}_{\text{GHZ}} = \frac{1}{2}\mathbb{1} - |\text{GHZ}\rangle\langle\text{GHZ}| \quad (9)$$

and $\text{Tr}(\mathcal{W}_{\text{GHZ}}|\text{GHZ}\rangle\langle\text{GHZ}|) = -E_{\sin^2}(|\text{GHZ}\rangle) = -1/2$. Similarly, for the states $|W\rangle \equiv (|001\rangle + |010\rangle + |100\rangle)/\sqrt{3}$ and $|\widetilde{W}\rangle \equiv (|110\rangle + |101\rangle + |011\rangle)/\sqrt{3}$ we have

$$\mathcal{W}_W = \frac{4}{9}\mathbb{1} - |W\rangle\langle W| \quad \text{and} \quad \mathcal{W}_{\widetilde{W}} = \frac{4}{9}\mathbb{1} - |\widetilde{W}\rangle\langle\widetilde{W}| \quad (10)$$

and $\text{Tr}(\mathcal{W}_W|W\rangle\langle W|) = -E_{\sin^2}(|W\rangle) = -5/9$, and similarly for $|\widetilde{W}\rangle$. For the four-qubit state $|\Psi\rangle \equiv (|0011\rangle + |0101\rangle + |0110\rangle + |1001\rangle + |1010\rangle + |1100\rangle)/\sqrt{6}$ the optimal witness is

$$\mathcal{W}_{\Psi} = \frac{3}{8}\mathbb{1} - |\Psi\rangle\langle\Psi| \quad (11)$$

and $\text{Tr}(\mathcal{W}_{\Psi}|\Psi\rangle\langle\Psi|) = -E_{\sin^2}(|\Psi\rangle) = -5/8$. In passing, we note that linear combinations of witnesses—preferably optimal ones—can be used to detect entanglement for mixed states, as we shall illustrate later. We also note that the non-optimal witnesses can also be of use, e.g., in classifying and detecting distinct types of entangled states; see Ref. [7]. Furthermore, as entanglement witnesses are *Hermitian* operators, they can, at least in principle, be realized experimentally.

Mixed states: We conclude by briefly commenting on the EW/GME connection for mixed states. The GME can be generalized to mixed states ρ via the *convex hull* construction (indicated by “co”). The essence of this construction is a minimization over all decompositions $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ into pure states:

$$E(\rho) \equiv (\text{co}E_{\sin^2})(\rho) \equiv \min_{\{p_i, \psi_i\}} \sum_i p_i E_{\sin^2}(|\psi_i\rangle). \quad (12)$$




FIG. 1: Contour plot of the detector, $\text{Tr}(\mathcal{W}(y)\rho(x))$. The darker the region, the more negative the detector. White regions correspond to positive values of the detector, where $\mathcal{W}$ is not a good entanglement witness, and are therefore left ungraded.

By using the analysis for pure states given above, we can rewrite the mixed-state entanglement as follows:

$$\begin{aligned} E(\rho) &\equiv \min_{\{p_i, \psi_i\}} \sum_i p_i E_{\sin^2}(|\psi_i\rangle) \\ &= -\max_{\{p_i, \psi_i\}} \sum_i p_i \min_{\mathcal{W}_i} \text{Tr}(\mathcal{W}_i |\psi_i\rangle\langle\psi_i|) \end{aligned} \quad (13a)$$

$$= -\max_{\{p_i, \psi_i\}} \sum_i p_i \text{Tr}(\mathcal{W}_{\psi_i} |\psi_i\rangle\langle\psi_i|), \quad (13b)$$

where $\mathcal{W}_{\psi_i}$ is the optimal EW corresponding to $|\psi_i\rangle$. Said equivalently, one can express the entanglement of ρ in terms of the optimal witnesses for the pure states that feature in the optimal decomposition. The GME for a mixed state, then, detects the minimum average of the “fidelities” between the pure components and their optimal witnesses. Although, for any entangled mixed state ρ , there always exists an entanglement witness $\mathcal{W}$, and by a trivial rescaling one can then have $\text{Tr}(\mathcal{W}\rho) = -E(\rho)$, it would be preferable to have a single, simple entanglement witness that is directly connected to the entanglement of the mixed state ρ . However, we do not know of any simple one.

Before ending our discussion of mixed states, we elaborate on a point made earlier, i.e., that linear combinations (with non-negative coefficients) of optimal entanglement witnesses can be used to establish entanglement of mixed states. For illustration, consider the following family of mixed states:

$$\rho(x) \equiv x|W\rangle\langle W| + (1-x)|\widetilde{W}\rangle\langle\widetilde{W}|, \quad (14)$$

the GME of which is calculated analytically in Ref. [5] and which is entangled for all values of $x \in [0, 1]$. We can actually construct EW’s that establish the entanglement of $\rho(x)$. Consider a linear combination of optimal witnesses of $|W\rangle$ and $|\widetilde{W}\rangle$:

$$\mathcal{W}(y) \equiv y \mathcal{W}_W + (1-y) \mathcal{W}_{\widetilde{W}}, \quad (15)$$

with $y \in [0, 1]$. If, for any given x , there exists a value of $y \in [0, 1]$ such that $\text{Tr}(\mathcal{W}(y)\rho(x)) < 0$ then $\rho(x)$ is evidently entangled. Figure 1 shows that this is indeed the case (see captions for details). This illustrates the usefulness of linear combinations of pure-state optimal witnesses. It would be interesting to know whether witnesses for bound entangled states be constructed by this approach.

Concluding remarks: Although the observations we have made are, from a technical standpoint, elementary, we nevertheless find it intriguing that two distinct aspects of entanglement—the geometric measure of entanglement

and entanglement witnesses—are so closely related. Furthermore, this connection sheds new light on the content of the geometric measure of entanglement. In particular, as entanglement witnesses are Hermitian operators, they can, at least in principle, be realized experimentally. Their connection with the geometric measure of entanglement ensures that the geometric measure of entanglement can, at least in principle, be verified experimentally.

Acknowledgments: This work was supported by NSF Award EIA01-21568 and by DOE DMS Award DEFG02-91ER45439 through the FS-MRL at UIUC. TCW acknowledges a Mavis Memorial Fund Scholarship.

-
- [1] See, e.g., M. Nielsen and I. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, 2000).
 - [2] For a review, see M. Horodecki, Quant. Info. Comp. **1**, 3 (2001), and references therein.
 - [3] A. Shimony, Ann. N.Y. Acad. Sci. **755**, 675 (1995).
 - [4] H. Barnum and N. Linden, J. Phys. A: Math. Gen. **34**, 6787 (2001).
 - [5] T. C. Wei and P. M. Goldbart, quant-ph/0212030.
 - [6] M. Lewenstein, B. Kraus, J. I. Cirac, and P. Horodecki, Phys. Rev. A **62**, 052310 (2000).
 - [7] A. Acín, D. Bruß, M. Lewenstein, and A. Sanpera, Phys. Rev. Lett. **87**, 040401 (2001).